REPORT DOCUMENTATION PAGE

Form Approved OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget. Paperwork Reduction Project (0704-0188), Washington, DC 20503.

1. AGENCY USE ONLY (Leave Blank)	2. REPORT DATE	3. REPORT TYPE AND DATES COVE	RED
	See Title Page	1 February 1987 to 31 Januar	y 1991
4. TITLE AND SUBTITLE			5. FUNDING NUMBERS
Use Title on Reprint One Copy	of Each Submitted		N00014-87-C-0146 0R0A444C-
6. AUTHOR(S)			43051 43051S4
See Individual Articles		BEIG	4303134
7. PERFORMING ORGANIZATION NAME	(S) AND ADDRESS(ES)	DECTERNI	8. PERFORMING ORGANIZATION
Vanderbilt University Nashville TN 37332 (615-322-2786)		ELECTE JAN 1 9 1996	REPORT NUMBER
9. SPONSORING/MONITORING AGENC	/ NAME(S) AND ADDRESS(ES)	G	10. SPONSORING/MONITORING AGENCY REPORT NUMBER
Office of Naval Research 800 North Quincy Street Arlington, VA 22217-5660			AGENOT REPORT NOMBER
11. SUPPLEMENTARY NOTES			
Each Paper Summarized on first pag	e. Journal articles submitted as contract	reports. All work performed under Gover	nment contract.
12a. DISTRIBUTION/AVAILABILITY STAT	EMENT		12b. DISTRIBUTION CODE
Approved for public release; dis	stribution unlimited.		
	Accesion For		
	NTIS CRA&I DTIC TAB Unannounced Justification		
	Ву		
	Distribution/		
	Availability Codes		
	Dist Avail and or Special	1996010	4 061
14. SUBJECT TERMS			15. NUMBER OF PAGES
Free Electron Lasers Medic Biomedical Instrumentation	ine Biology		00 16. PRICE CODE
Energy Cells			
17. SECURITY CLASSIFICATION OF REPORT	18. SECURITY CLASSIFICATION OF THIS PAGE	19. SECURITY CLASSIFICATION OF ABSTRACT	20. LIMITATION OF ABSTRACT
UNCLASSIFIED	UNCLASSIFIED	UNCLASSIFIED	UL

NSN 7540-01-280-5500

Standard Form 298 (Rev. 2-89) Prescribed by ANSI Std 239-18 298-102

WENTER TO BE STOP

PHYSICS OF FREE-ELECTRON-LASER APPLICATIONS

Physics of Free-Electron-Laser Applications in the Visible and Infrared

Introduction		9/2
Two-color free-electron laser driven by a radio-frequency linear accelerator	H. A. Schwettman, T. I. Smith	973
Application of a two-color free-electron laser to condensed-matter molecular dynamics	Dana D. Dlott, Michael D. Fayer	977
Infrared free-electron laser as a probe of vibrational dynamics on surfaces	N. J. Tro, D. A. Arthur, S. M. George	995
Excitation spectroscopy of thin-film amorphous semiconductors using a free- electron laser	N. J. Ristein, B. Hooper, P. C. Taylor	1003
Effects of introducing a gas into the free-electron laser	R. H. Pantell, A. S. Fisher, J. Feinstein, A. H. Ho, M. Özcan, H. D. Dulman, M. B. Reid	1008
Optical spectroscopic diagnosis of cancer and normal breast tissues	R. R. Alfano, Asima Pradhan, G. C. Tang, S. J. Wahl	1015
Ultrafast carrier relaxation in hydrogenated amorphous silicon	P. M. Fauchet, D. Hulin	1024
Far-infrared photo-Hall experiments on GaAs:Si	J. Kaminski, J. Spector, C. T. Foxon, T. O. Klaassen, W. Th. Wenckebach	1030
Material applications of the far-infrared free-electron laser	John D. Simon, John E. Crowell, John H. Weare, David R. Miller	1035
Applications of far-infrared free-electron lasers to condensed-matter physics	W. M. Dennis	1045
Enhancement of ligand binding to myoglobin by far-infrared excitation from a free-electron laser	Bernard Gerstman, Mark Roberson, Robert Austin	1050
Proposal for the direct electromagnetic generation of coherent terahertz acoustic phonons in semiconductor superlattices at the University of California, Santa Barbara far-infrared free-electron-laser facility	T. E. Wilson	1058
Physics of Free-Electron-Laser Applications in the Vacuum Ultraviolet–Extreme Ultraviolet		
Introduction		1061
Fluorescence spectroscopy as a probe of the electronic structure and the dynamics of rare-gas clusters	T. Möller, G. Zimmerer	1062
Prospects for photoionization studies of weakly bound molecular complexes using free-electron-laser vacuum-ultraviolet radiation	E. A. Walters, J. R. Grover	107
Precision spectroscopy on stored, highly charged ions using free-electron-laser radiation	D. A. Church, S. D. Kravis	107
Free-electron lasers in ultraviolet photobiology	Thomas P. Coohill, John C. Sutherland	1079
Testing the photon-photon sector of quantum electrodynamics with free- electron lasers	W. Becker, J. K. McIver, R. R. Schlicher	108



Physics of Free-Electron-Laser Applications in the Visible and Infrared

INTRODUCTION

It is now eighteen years since John Madey published a paper pointing out that a high-brightness relativistic electron beam traversing a spatially periodic magnetic field could stimulate the emission of photons over a broad range of wavelengths, indeed, from the far infrared to the ultraviolet. In a way, the free-electron laser was the ultimate homage paid by the laser, viewed as an optical device, to its antecedents in radar and electron-beam science and technology dating back into the 1940's.

In the intervening years, successful infrared and visible free-electron-laser (FEL) experiments, for example, at Stanford, Orsay, Santa Barbara, and Los Alamos, have shown significant promise for applications based on the unique optical characteristics of the FEL. A variety of accelerators can provide the high-brightness electron beams necessary for the FEL: room-temperature pulsed linear accelerators, superconducting accelerators, storage rings, and Van de Graaff generators have all been successfully used so far for this purpose. The existence of this variegated collection of pumps for the stimulated emission generated in the FEL implies a correspondingly broad range of temporal pulse shapes, interpulse spacings, pulse-repetition frequencies, output powers, and spectral ranges for users.

With the increasing maturity of the free-electron laser comes a new phase of scientific opportunity for those who are primarily laser users rather than laser physicists. During the past two years, FEL users' facilities at Stanford University and the University of California at Santa Barbara began to provide significant quantities of time to photon users, particularly in surface and materials science and biomedical studies. In the coming year, new FEL users' facilities devoted to biomedical and materials research as well as to FEL development will begin to operate at Vanderbilt and Duke; plans for additional facilities of this type are already far advanced.

Free-electron lasers now support significant experimental activities in a wide range of scientific and technological applications, including biomedical research. The spectrum of wavelengths now demonstrated in FEL's ranges from 0.25 to $800~\mu m$, the region that includes atomic and molecular transitions as well as many elementary excitations in solids. While at least some parts of this spectral region can be covered by conventional lasers, the FEL has at least two major benefits compared with conventional lasers:

- The variety of FEL accelerator types offers the potential for decoupling the laser temporal-pulse structure and power output from the constraints of discharge or laser-pumping schemes and
- Beyond a wavelength of approximately 10 μ m, the FEL has significantly greater tunability and higher power than any currently available lasers.

In addition, the fact that the interpulse spacing can be set much shorter than that available from mode-locked pulse trains (the Mark III FEL at Stanford and slated for installation at Vanderbilt, for example, has a rf accelerator frequency of 2.865 GHz, so that its pulses are 330 psec apart) makes it possible to carry out experiments in which the laser pulses follow one another on a time scale comparable with many relaxation times of physical, chemical, and biological interest.

In this special issue of Journal of the Optical Society of America B, we have collected a round dozen papers describing the physics of FEL applications in the visible and infrared. First, the companion papers on two-color FEL technology and applications suggest new opportunites in fundamental science that can be attained with the unique technology of the free-electron laser. In the near infrared, additional condensed-matter studies and surface scienceparticularly vibrational spectroscopy—have become possible with existing free-electron lasers. The gas-loaded freeelectron laser can shift infrared FEL output into the visible. where work done with existing tabletop lasers can suggest ways in which the FEL, with its unique temporal pulse characteristics, might enhance our understanding of laserinduced processes both in materials science and in biomedical research. Finally, the far-infrared region was pinpointed early on as a place where exciting studies could be carried out in condensed-matter physics and biological physics. The papers submitted for this issue covering both recent research applications and new possibilities for direct excitation of far-infrared vibrational modes in solids suggest that this promise of the FEL is already being realized.

He tree who who was a company to the company to the

he

onc

However, as with any new photon source, much of the challenge in developing new science will revolve around "thinking the unthinkable," or at least thinking the heretofore unthought. Thus the material ranges from reports of experimental results to inform speculations on possible future directions in FEL applications based on the unique properties of the FEL. Even though the Journal of the Optical Society of America B is an archival journal with a retrospective point of view, we hope that the speculation will stimulate as yet unthought, forward-looking applications of the FEL in the larger optical-science community.

Richard F. Haglund, Jr.
Howard Schlossberg
Feature Editors
Physics of Free-Electron-Laser Applications
in the Visible and Infrared